

RESEARCH ARTICLE

PHYSICAL ACTIVITY AS COPING STRATEGY FOR ACADEMIC STRESS AMONG UNDERGRADUATE FEMALE STUDENT

1. Director Physical Education. Bilquis Postgraduate College for Women, Rawalpindi, Pakistan
2. Senior Lecturer. College of Physical Therapy. Northwest Institute of Health Sciences. Peshawar. Pakistan.
3. Physical Education Expert. Pakistan Sports Board. Islamabad Pakistan.

Correspondence Hamsa Ashraf

Director Physical Education. Bilquis Postgraduate College for Women PAF, Nur Khan Base. Rawalpindi, Pakistan
E-mail: ayat_a2@yahoo.co.uk

Received on: 31-01-2020

Revision on: 01-05-2020

Published on: 30-06-2020

Citation; Ashraf H, Hussain SA, Haseeb M, Haider HMM. Physical activity as coping strategy for academic stress among undergraduate female student. T Rehabil. J. 2020;04(01):144-148. doi: [21-2017/re-trjvol04iss01p144](https://doi.org/10.21017/re-trjvol04iss01p144)

Hamsa Ashraf¹: Conception & designing, analysis, writing; revised and accountable for all aspects
Syed Alamdar Hussain²: Analysis & interpretation of data, writing; revised and accountable for all aspects.
Musa Haseeb³: writing; revised and accountable for all aspects.

ABSTRACT

Objective: To determine the effectiveness of physical activity as a coping strategy to reduce academic stress among undergraduate female students. **Methodology:** A Randomized control trail (NCT04221022) was conducted in Bilquis Postgraduate College for Women PAF, Nur Khan Base, Rawalpindi. A total of n=37 undergraduate inactive female (>1-month) students, age between 18-24 years, with moderate level of academic stress measured on academic stress scale were included in the study. While students with diagnosed psychological disorders, systematic diseases were excluded. The participants randomly divided into Light Physical Activity (LPA), Moderate Physical Activity (MPA) and Vigorous Physical activity (VPA) groups. The academic stress scale (ASS) was used to observe level of academic stress among participant at baseline and after six weeks of intervention. **Results:** The mean age of female study participants was 20.24±1.44 years. Pre-Post analysis showed that all group significantly improve ($p<0.001$) level of academic stress after 6 week intervention. The academic stress was significantly improved in moderate PA group as compare to vigorous PA (15.17±12.39 Ver. 44.62±36.94, $p=0.019$). **Conclusion:** All types of physical activities were effective as a coping strategy for reducing academic stress among female undergraduate students. But moderate level of physical activity like brisk walking was significantly associated with reduction in academic stress as compare to vigorous and light physical activities.

Keywords: Cognition, mental health, physical activity, stress.

INTRODUCTION

Physical activity (PA) is defined as any bodily movements, which are produced by skeletal muscles that results in energy expenditure and classified as light, moderate and vigorous physical activity. PA is effective in preventing and treating high blood pressures, diabetes, osteoporosis and depression. People who performs PA are more efficient and healthier.¹ Whereas risk of developing hypertension, coronary heart disease, diabetes and obesity increases due to physical inactivity.²

Now-a-days stress is a typical chunk of each individual's life.³ Excessive work leads to the feeling of physical stress whereas improper daily life routine cause mental stress.⁴ Stressors such as academic, self-imposed, economic, and health related may affect the college students.⁵ The education system is quite stressful these days. A student's life and specifically a female student is subjected to diverse kinds of stressors such as the burden of academics, grades, undefined future and family problems.⁶ stress also leads to the issues of concentration, tension, anxiety, focus and attention.⁷

Moreover, students are also overloaded with quizzes, assignments, presentations and exams and all these factors are working as stressors in students' life which may effects the academic achievements.⁸ Academic stress is mental distress

which may occurs due to the pressure of educational failure, examination, class participation, competing other class mates and extra-curricular activities.^{9,10} Due to which students face emotional, physical, and social problems which might affect their learning capability and academic performance.⁶

Regular PA is effective for health life style and it is suggested as a coping strategy for stress.¹¹ In some previous studies, it has been demonstrated that most of the students do not participate in physical activity, whereas engaging in it can be a potent tool to enhance the physiological wellbeing of the student and minimizing there academic stress.^{12,13} However, in literature there is also limited evidence that physical activity is being properly used as an effective therapy in the management of academic stress. This study was conducted to determine the effectiveness of physical activity as a coping strategy to reduce academic stress among female participants.

METHODOLOGY

A randomized clinical trial (NCT04221022) was conducted in Bilquis Postgraduate College for Women PAF Nur Khan Base Rawalpindi after the permission from Principal. A total of n=37 undergraduate inactive female (>1-month)

students, age between 18-24 years, with moderate level of academic stress measured on academic stress scale were included in the study. While students with diagnosed psychological disorders, systematic diseases were excluded from the study.

The participants were randomly divided into three groups, through the sealed envelope method such as Light Physical Activity (LPA), Moderate Physical Activity (MPA) and Vigorous Physical activity (VPA) group. (Figure 1)




Figure 1: CONSORT Diagram

The guidelines regarding physical activity for intervention were obtained from Rapid Assessment of Physical Activity (RAPA) questionnaire which has constructed validity and reliability.¹⁵ The duration of study were 6 weeks and each group receiving intervention for 5 days/week. The participants were ask to perform Leisure Walk for 35 minutes in LPA group, Brisk Walk for 30 minutes in MPA group and jogging for 15 minutes in VPA group. The information regarding age, BMI and semester was obtained at baseline. The academic stress scale (ASS) was used to observe level of academic stress among participant at baseline and after six weeks, which is also a valid and reliable tool.¹⁶ The assumptions of parametric tests were met, so for within group changes, paired sample t-test and for comparison One Way ANOVA with Tukey HSD post hoc was used. The level of significance was set as

$p < 0.05$, and SPSS ver. 21 was used to analyse the data.

RESULTS

The mean age of female study participants was 20.24 ± 1.44 years and mean BMI $20.23 \pm 0.26 \text{ kg/m}^2$, which showed that majority of participant, had normal BMI score. Majority of students were in 2nd (n=10) and 4th (n=14) semester respectively (Figure 2).

Pre-Post analysis showed that all group showed improvement in level of academic stress after 6 week intervention, while comparison between groups showed significant difference ($p = 0.024$) among the groups as shown in figure 3.




Figure 2: Semester wise distribution of participants




Figure 3: With-in group changes in ASS

Post Hoc analysis with Tukey HSD showed that the academic stress was significantly improved in moderate PA group as compare to vigorous PA (15.17±12.39 Ver. 44.62±36.94, $p=0.019$). But no significant difference was observed between

moderate and light PA group (15.17±12.39 Ver. 33.5±19.71, $p=0.196$) as well as light and vigorous PA group (33.5±19.71 Ver. 44.62±36.94, $p=0.543$) as shown in Figure 4.




Figure 4: Comparison of ASS among LPA, MPA & VPA groups

DISCUSSION

According to the results, each level of physical activity reduced the academic stress level at the end of 6 weeks but moderate level of physical activity e.g. brisk walk significantly improve academic stress symptoms as compare to light and vigorous physical activity.

In this study, it was concluded that moderate level of PA was significantly associated with the academic stress, which correlates with the previous study conducted by Ulrich A. In which it has been demonstrated that moderate level of PA i.e. 150 minutes/week is effective for reducing stress and improve health in undergraduate students.¹⁷

Moreover, it has also been determined that regular moderate and vigorous intensity PA improves mental and cognitive health.¹⁸ A study conducted by James A. Blumenthal et al. discussed that 12 weeks of aerobics and strength training is effective for reduction in mental stress.¹⁹ However, according to the results of this study, brisk walking for a time period of 6 weeks can significantly use as a coping strategy for mental stress. This occurs due to the cortisol level, which is considered as stress hormone. Cortisol is released during physical activity such as brisk walking which reduces the stress level while secretion of cortisol is decreased or inhibited due to low PA.²⁰

Additionally, another possible mechanism is the increased concentration of serotonin and other neurotransmitters due to PA, which are associated with the endorphin effects and decrease the negative effects of stress.²¹ This supports the current findings of the study that moderate intensity of physical activity for 6 weeks reduces the stress level and improves mental health of college students. And stress is the leading cause of poor academic performance in college students.¹⁴

Moreover, in the current study no significant difference has been observed in the light and vigorous PA after 6 weeks of intervention. However, in a previous literature, it has been discussed that vigorous PA is inversely associated with the stress level among college students.²² On the other hand, walking in nature is also effective for the mental health.²³ However, there's less evidence available for the leisure walk and the academic stress in college students.

CONCLUSION

All type of physical activities were effective for coping up with the academic stress among female undergraduate students. But moderate level of physical activity like brisk walking was significantly associated with the academic stress, as compare to vigorous and light physical activities. The current study was single centre study as well as no confounding variables like age socioeconomic status were included in the analysis. So multi-centre study, while focusing confounding variables, is recommended for future research.

REFERENCES

1. Abou Elmagd M. Benefits, need and importance of daily exercise. *Int. J. Phys. Educ. Sports Health.* 2016;3(5): 22-27.
2. Powell KE, Blair SN. The public health burdens of sedentary living habits: theoretical but realistic estimates. *Med. Sci. Sports Exerc.* 1994;26(7):851-6.
3. Ruscio AM, Gentes EL, Jones JD, Hallion LS, Coleman ES, Swendsen J. Rumination predicts heightened responding to stressful life events in major depressive disorder and generalized anxiety disorder. *J. Abnorm. Psychol.* 2015;124(1):17.
4. Kofoworola OH, Alayode AM. Strategies for managing stress for optimal job performance. *Int. J. Psychol. Stud.* 2012;4(2):162.
5. SAJID MB, HAMID S, SABIH F, SAJID A. Stress and coping mechanisms among college students. *J. Pak. Psychiatr. Soc.* 2017;14(3):31-4.
6. Shaikh B, Kahloon A, Kazmi M, Khalid H, Nawaz K, Khan N, et al. Students, stress and coping strategies: a case of Pakistani medical school. *Educ. Health.* 2004;17(3):346-53.
7. Cruz SY, Fabian C, Pagán I, Ríos JL, González AM, Betancourt J, et al. Physical activity and its associations with sociodemographic characteristics, dietary patterns, and perceived academic stress in students attending college in Puerto Rico. *P. R. Health. Sci. J.* 2013;32(1).
8. Chi T. Physical activity is defined as "any bodily movement produced by skeletal muscles that results in caloric expenditure"(Pender, 1996, p. 185). Definitions of exercise are complex and vary according to scientific. *Complementary Alternative Therapies in Nursing.* 2009:349.
9. Morrison V, Bennett P. *An introduction to health psychology:* Pearson Education; 2009.
10. Ströhle A. Physical activity, exercise, depression and anxiety disorders. *J. Neural. Transm.* 2009;116(6):777.
11. Wijndaele K, Matton L, Duvigneaud N, Lefevre J, De Bourdeaudhuij I, Duquet W, et al. Association between leisure time physical activity and stress, social support and coping: A cluster-analytical approach. *Sport Exerc.* 2007;8(4):425-40.
12. Nguyen-Michel ST, Unger JB, Hamilton J, Spruijt-Metz D. Associations between physical activity and perceived stress/hassles in college students. *Stress Health: Stress Health.* 2006;22(3):179-88.
13. Stephens T. Physical activity and mental health in the United States and Canada: evidence from four population surveys. *Prev. Med.* 1988;17(1):35-47.
14. Lawrence NS, Aduka SZ, Desa ZM. Factors of Stress Affecting the Academic Performance among the University Students.

15. Aktan R, Ozalevli S, Yalcinkaya G, Alpaydin AO, Sevinc C. Validation of the rapid assessment of physical activity questionnaire (RAPA) in COPD patients. *Eur Respiratory Soc*; 2018.
16. Bedewy D, Gabriel A. Examining perceptions of academic stress and its sources among university students: The Perception of Academic Stress Scale. *Health Psychol. Open*. 2015;2(2):2055102915596714.
17. Ulrich A. The Impact of Physical Activity on Stress. *Methods for Stress Management*. 2017.
18. Nakagawa T, Koan I, Chen C, Matsubara T, Hagiwara K, Lei H, et al. Regular Moderate-to Vigorous-Intensity Physical Activity Rather Than Walking Is Associated with Enhanced Cognitive Functions and Mental Health in Young Adults. *Int. J. Environ. Res. Public Health*. 2020;17(2):614.
19. Blumenthal JA, Fredrikson M, Kuhn CM, Ulmer RL, Walsh-Riddle M, Appelbaum M. Aerobic exercise reduces levels of cardiovascular and sympathoadrenal responses to mental stress in subjects without prior evidence of myocardial ischemia. *Am. J. Cardiol*. 1990;65(1):93-8.
20. Jin P. Efficacy of Tai Chi, brisk walking, meditation, and reading in reducing mental and emotional stress. *J. Psychosom. Res*. 1992;36(4):361-70.
21. Wu X, Tao S, Zhang Y, Zhang S, Tao F. Low physical activity and high screen time can increase the risks of mental health problems and poor sleep quality among Chinese college students. *PLoS ONE*. 2015;10(3):e0119607.
22. VanKim NA, Nelson TF. Vigorous physical activity, mental health, perceived stress, and socializing among college students. *Am. J. Health Promot*. 2013;28(1):7-15.
23. Marselle MR, Warber SL, Irvine KN. Growing resilience through interaction with nature: Can group walks in nature buffer the effects of stressful life events on mental health? *Int. J. Environ. Res. Public Health*. 2019;16(6):986.

Disclaimer: None to declare.

Conflict of Interest: None to declare.

Funding Sources: None to declare.